## No XVII.—THE AFFINITIES OF ERYX JACULUS AND ERYX JOHNI.

Captain Ingoldby's suggestion in our last Journal (Vol. XXIX, p. 127) that Eryx johni does not deserve recognition as a species apart from E. jaculus has prompted me to refer to my notes on the specimens I have examined of both supposed species. I have for some years considered the two so closely allied that I have been guided largely by the locality of a specimen in identifying the species. In this Journal (Vol. XIX, p. 1,000). I identified a specimen from Jhelum as jaculus, the lepidosis being nearer to the description of this species in Boulenger's Catalogue (Vol. 1, p. 125) than that of johni (loc. sit. p. 127).

I think my records given below giving details of the lepidosis upon which the two supposed species are separated, strongly support Captain Ingoldby's

view that johni should be rejected as a species apart from jaculus.

	Costals.							
Serial No.	Two head-lengths behind head.	Midbody.	Two head-lengths before vent.	Ventrals.	Subcaudals.	Locality.		
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22	51 48 42 47 44 47 45 48 46 48 46 48 47 42 44 45 49 46 46 47 47 47 54 48	57 53 49 52 50 52 51 54 52 49 52 53 47 51 53 56 51 52 54 52 61 52	37 37 37 35 35 39 37 39 37 ? 38 41 34 38 42 37 36 41 40 46 40	190 190 190 198 195 197? 190 199 191 193 194? 190? 199 200 195 200 193 189 192 193	26 24 33 27 30 26 28 25 26 26 26 26 26 33 33 29 35 33 37 30 30 37	Iraq. (Basra.) Do. do. Do. do. Muhammerah, S Persia, Do.		
$\begin{array}{c} 23 \\ 24 \end{array}$	47 39	52 48	41 53	193 192	29 34	Do. ( do. ). Do. ( do. ).		
25	51	59	42	201	31	Do. ( do. ).		
26	49	58	42	198	34	Do. ( do. ).		
27	54	60	41	215	30	Do. ( do. ).		
28	52	62	45	213	••	S. India (Bangalore)		
	1			3				

Colouration.—Captain Ingoldby says in the few young Indian specimens he has seen the body was quite unmarked. In young Sind specimens, however, and up to the time they are half grown, one sees ill-defined black bars on the body, more pronounced posteriorly.

I have never seen any specimens from Iraq and Persia that had acquired the uniform dark brown dorsal colouration so typical of the adult Indian form of the species, but immature Indian specimens resemble in colouration the speci

mens so typical of Iraq and Persia.

Dentition. - I have skulls of six Indian specimens that I have hitherto labelled johni, and two from Persia and Iraq identified as jaculus. A study of the teeth supports the view that they constitute a single species as will be seen from the following details.

	Maxillary.			Palatine.	Pterygoid.	Mandibul- ar.	Locality.
Left		11	1	4	3	13	
Right		11		4	3	13	Trichinopoly.
Left		10		4	5	12	
Right		10		4	4	12	Karachi.
Left		10		4	4	?	
Right		10		4	4	?	Karachi.
Left		10		3	4	11	
Right		10		3	4	12	Jhelum.
Left		10		3	4	12	
Right		10		3	4	12	Jhelum.
Left		10		4	4	13	
Right		10		4	4	13	Jhelum.
Left		11		3	4	14	
Right		10		3	4	16	Muhammerah.
Left		10		3	4	11	
Right	• •	9		3	4	11	Basra.
			1				

The maxillary series are anododont, syncranterian and scaphiodont. The palatine anododont, isodont. The pterygoid anododont, isodont. The mandi-

bular anododont, strongly scaphiodont.

Distribution. The distribution of this snake is interesting as only one other species, viz., Echis carinatus occurring in South India extends to Persia and Iraq.

KARACHI, 4th August 1923.

F. WALL, COLONEL, I.M.S.

## No. XVIII—AN UNUSUALLY LARGE LABEO FROM THE INDRAYANI RIVER.

The following incident, which may be of interest to fishermen in these parts

occurred yesterday.

In company with Major and Mrs. T. W. Corbett, I had been out after green pigeon and was returning to Poona via the Nasik Road. On reaching the causeway over the river Indrayani we stopped for a smoke and to give the cars a drink. The water on the upstream side of the causeway was about a foot below road level and about five or six feet below the road on the lower side, with a series of small waterfalls rushing out from the sluices in the middle, the